

Cardiometabolic Statistics

More than
1 in 3
US adults have
metabolic syndrome.¹



Woman with metabolic syndrome may be at

62%




greater risk for infertility.²

Non-alcoholic fatty liver disease (NAFLD) is the

most common cause

of chronic liver disease.³



Over 30%

of adults in the United states have Non-alcoholic fatty liver disease (NAFLD).⁴

Statistics show that statins are only effective in preventing a heart attack in
1 out of 100
people.⁵



Statin use increased 10% between 2003 and 2013 for US adults over 40 years old.⁶

Over 20%

of youths aged 6-19 have at least 1 abnormal cholesterol measure.⁷


Over 25%

of adults have hypertension, up to half of those may be uncontrolled hypertension.⁸

One analysis found that

Over 60%

of individuals with bipolar disorder or schizophrenia have cardiometabolic comorbidities.⁹



References

1. Moore JX, Chaudhary N, Akinyemiju T. Metabolic Syndrome Prevalence by Race/Ethnicity and Sex in the United States: 2017. National Health and Nutrition Examination Survey, 1988–2012. *Prev Chronic Dis* 2017;14:160287. DOI: <http://dx.doi.org/10.5888/pcd14.160287>.
2. Perumpail BJ, Khan MA, Yoo ER, Cholankeril G, Kim D, Ahmed A. Clinical epidemiology and disease burden of nonalcoholic fatty liver disease. *World J Gastroenterol*. 2017;23(47):8263–8276. doi:10.3748/wjg.v23.i47.8263
3. Grieger JA, Grzeskowiak LE, Smithers LG, Bianco-Miotto T, Leemaqz SY, Andraweera P, Poston L, McCowan LM, Kenny LC, Myers J, Walker JJ, Norman RJ, Dekker RJ, Roberts CT. Metabolic syndrome and time to pregnancy: a retrospective study of nulliparous women. *BJOG*. 2019 Jun;126(7):852–862. doi: 10.1111/1471-0528.15647.
4. Kennedy-Martin T, Bae JP, Paczkowski R, Freeman E. Health-related quality of life burden of non alcoholic steatohepatitis: a robust pragmatic literature review. *J Patient Rep Outcomes*. 2018;2:28. Published 2018 Jun 19. doi:10.1186/s41687-018-0052-7.
5. David M Diamond, Uffe Ravnskov. How statistical deception created the appearance that statins are safe and effective in primary and secondary prevention of cardiovascular disease. *Expert Review of Clinical Pharmacology*, 2015; 8 (2): 201 DOI: 10.1586/17512433.2015.1012494.
6. Salami JA, Warraich H, Valero-Elizondo J, Spatz ES, Desai NR, Rana JS, Virani SS, Blankstein R, Khera A, Blaha MJ, Blumental RS, Lloyd-Jones D, Nasir K. National Trends in Statin Use and Expenditures in the US Adult Population From 2002 to 2013. *Insights From the Medical Expenditure Panel Survey*. *JAMA Cardiol*. 2017;2(1):56–65. doi:10.1001/jamacardio.2016.4700.
7. Benjamin EJ, Virani SS, Callaway CW, et al. American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. Heart Disease and Stroke Statistics-2018 Update: A Report From the American Heart Association. *Circulation*. 2018 Mar 20;137(12):e67–e492. doi: 10.1161/CIR.0000000000000558.
8. Table 53. Selected health conditions and risk factors, by age: United States, selected years 1988–1994 through 2015–2016. 2017 Health, United States. <https://www.cdc.gov/nchs/data/hsr/2017/053.pdf>. Last accessed 09/14/19.
9. Correll CU, Ng-Mak DS, Stafkey-Mailey D, Farrelly E, Rajagopalan K, Loebel A. Cardiometabolic comorbidities, readmissions, and costs in schizophrenia and bipolar disorder: a real-world analysis. *Ann Gen Psychiatry*. 2017 Feb 10;16:9. doi: 10.1186/s12991-017-0133-7.



THE INSTITUTE FOR
FUNCTIONAL
MEDICINE®